

University of Massachusetts Medical School

eScholarship@UMMS

---

PEER Liberia Project

UMass Medical School Collaborations in Liberia

---

2021-04-22

## Acute Stroke Imaging

Jose Gavito-Higuera  
*RAD-AID*

Let us know how access to this document benefits you.

Follow this and additional works at: [https://escholarship.umassmed.edu/liberia\\_peer](https://escholarship.umassmed.edu/liberia_peer)



Part of the [Cardiovascular Diseases Commons](#), [Diagnosis Commons](#), [Family Medicine Commons](#), [Medical Education Commons](#), [Nervous System Diseases Commons](#), and the [Radiology Commons](#)


---

### Repository Citation

Gavito-Higuera J. (2021). Acute Stroke Imaging. PEER Liberia Project. <https://doi.org/10.13028/kdpq-he34>. Retrieved from [https://escholarship.umassmed.edu/liberia\\_peer/75](https://escholarship.umassmed.edu/liberia_peer/75)

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in PEER Liberia Project by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

# Acute Stroke Imaging



TEXAS TECH UNIVERSITY  
HEALTH SCIENCES CENTER  
EL PASO

## Jose Gavito-Higuera, MD

Assistant Professor, Neuroradiology  
Texas Tech University Health Sciences Center  
University Medical Center of El Paso  
El Paso, Texas  
[jose.gavito@ttuhsc.edu](mailto:jose.gavito@ttuhsc.edu)

1

## Overview:

- I. Acute stroke imaging algorithm
- II. CT / CTA / CTP Stroke assessment
- III. Review Cases

2

## “Time is Brain”



- 1.9 million neurons die each minute in which stroke is untreated
- But it depends on if you are a slow, moderate or fast progressor and this may be dependent on individual collaterals

3

## “Time is Brain”



Success of acute stroke intervention depends on :

- Patient selection
- Symptom onset to treatment time

4

# Acute Stroke – Clinical Evaluation

- Neurological exam
  - National Institutes of Health Stroke Scale

- Time of symptom onset
  - < 3 – 4.5 hrs IV tPA
  - < 6 hrs IA therapy

ECASS III study  
Hacke W, At al NEJM, 2008

- tPA inclusion / exclusion
  - ACEP/AAN (2013)

5

- National Institutes of Health Stroke Scale (NIHSS)

## Score:

0 : No Stroke

1- 4 : Minor Stroke

5 – 15 : Moderate Stroke

16 – 20 : Moderate to Severe

21 – 42 : Severe Stroke

National Institutes of Health Stroke Scale	
Score = 0 No stroke	Score = 5-15 Moderate stroke
Score = 1-4 Minor stroke	Score = 16-20 Moderate to severe stroke
	Score = 21-42 Severe stroke
National Institutes of Health Stroke Scale score	
1a. Level of consciousness	0 = Alert; keenly responsive 1 = Not alert, but arousable by minor stimulation 2 = Not alert; requires repeated stimulation 3 = Unresponsive or responds only with reflex
1b. Level of consciousness questions:	0 = Answers two questions correctly 1 = Answers one question correctly 2 = Answers neither question correctly
What is the month?	
What is your age?	
1c. Level of consciousness commands:	0 = Performs both tasks correctly 1 = Performs one task correctly 2 = Performs neither task correctly
Open and close your eyes.	
Grip and release your hand.	
2. Best gaze	0 = Normal 1 = Partial gaze palsy 2 = Forced deviation
3. Visual	0 = No visual loss 1 = Partial hemianopia 2 = Complete hemianopia 3 = Bilateral hemianopia
4. Facial palsy	0 = Normal symmetric movements 1 = Minor paralysis 2 = Partial paralysis 3 = Complete paralysis of one or both sides
5. Motor arm	0 = No drift 1 = Drift 2 = Some effort against gravity 3 = No effort against gravity; limb falls 4 = No movement
5a. Left arm	
5b. Right arm	
6. Motor leg	0 = No drift 1 = Drift 2 = Some effort against gravity 3 = No effort against gravity 4 = No movement
6a. Left leg	
6b. Right leg	
7. Limb ataxia	0 = Absent 1 = Present in one limb 2 = Present in two limbs
8. Sensory	0 = Normal; no sensory loss 1 = Mild-to-moderate sensory loss 2 = Severe to total sensory loss
9. Best language	0 = No aphasia; normal 1 = Mild to moderate aphasia 2 = Severe aphasia 3 = Mute, global aphasia
10. Dysarthria	0 = Normal 1 = Mild to moderate dysarthria 2 = Severe dysarthria
11. Extinction and inattention	0 = No abnormality 1 = Visual, tactile, auditory, spatial, or personal inattention 2 = Profound hemi-inattention or extinction
Total score = 0-42.	

Eleven-category assessment, assigning different point value to different function

6

## Acute Stroke – Clinical Evaluation

- Neurological exam
  - National Institutes of Health Stroke Scale

- Time of symptom onset
  - < 3 – 4.5 hrs IV tPA
  - < 6 hrs IA therapy


ECASS III study  
Hacke W, et al NEJM, 2008

- tPA inclusion / exclusion
  - ACEP/AAN (2013)

The other clinical component is the time of stroke. There are multiple studies (tPA has benefit but only in a certain time of window).


7

## Stroke Activation Protocol




8

# Stroke Activation Protocol




9

# Acute Stroke – Imaging Evaluation




10

## Acute Stroke – Imaging Evaluation




11

## Acute Stroke – Imaging Evaluation




12

## Acute Stroke – Imaging Evaluation




13



14




## Acute Stroke – Imaging Evaluation




15

## Acute Stroke – Imaging Evaluation




16

## Acute Stroke – Imaging Evaluation




17

## Acute Stroke – Imaging Evaluation




18

## Acute Stroke – Imaging Evaluation




19


## Acute Stroke – Imaging Evaluation



20



21



22

## CT signs of ACUTE stroke:



- Hypoattenuation
- Loss of gray white matter differentiation
- Sulcal effacement and mass effect
- Hyperdense vessel sign (MCA, dot-M2)

23

## Loss of gray white matter differentiation

- Basal ganglia sign



- Insular ribbon sign



24

## CT signs of ACUTE stroke:




- Hypoattenuation
- Loss of gray white matter differentiation
- Sulcal effacement and mass effect
- Hyperdense vessel sign (MCA, dot-M2)

25

## Correlating finding?

- Dense MCA sign



26


## Case 1:



What CT acute stroke finding shown?

- a) Insular ribbon sign
- b) Loss of gray white matter differentiation
- c) Dense vessel sign
- d) Hemorrhagic conversion


27



28

## CT stroke Volume

- Alberta stroke program early CT score (ASPECTS)



The diagram illustrates the ASPECTS Score system, which evaluates 10 regions of the MCA territory on axial CT scans. The regions are: C (Caudate), I (Insular ribbon), IC (Internal Capsule), L (Lentiform nucleus), M1 (Anterior MCA cortex), M2 (MCA cortex lateral to the insular ribbon), M3 (Posterior MCA cortex), M4 (Anterior MCA superior territory), M5 (Lateral MCA superior territory), and M6 (Posterior MCA superior territory). Each region is represented by a checkbox in the legend. The ASPECTS Score is calculated as the number of regions without evidence of stroke, out of a total of 10.

**ASPECTS Score**

- ☐ C- Caudate
- ☐ I- Insular ribbon
- ☐ IC- Internal Capsule
- ☐ L- Lentiform nucleus
- ☐ M1- Anterior MCA cortex
- ☐ M2- MCA cortex lateral to the insular ribbon
- ☐ M3- Posterior MCA cortex
- ☐ M4- Anterior MCA superior territory
- ☐ M5- Lateral MCA superior territory
- ☐ M6- Posterior MCA superior territory


ASPECTS Score = /10

**Practical points:**

- Total 10 MCA territory allotted where each area equals to 1 point
- Normal CT scan receive a score of 10 points
- A score of  $<7$  is equal to one-third of the MCA territory
- A score of zero indicates diffuse ischemia involvement throughout the MCA territory

29

## Case 2:



Two axial CT scan images are shown side-by-side. The left image is a normal CT scan with a score of 10. The right image shows a large area of hypodensity in the right hemisphere, indicating a large stroke. The ASPECTS score for the right image is 5.


**ASPECTS score?**

- a) 10
- b) 9
- c) 8
- d) 7
- e) 6
- f) 5

30



### Case 3:




ASPECTS score?

- a) 7
- b) 6
- c) 5
- d) 4
- e) 3
- f) 2

31

### Case 4:




ASPECTS score?

- a) 10
- b) 9
- c) 8
- d) 7
- e) 6
- f) 5

32


## CT stroke Volume

- Alberta stroke program early CT score (ASPECTS)



33

## Question 1:




Which of the following most likely makes a patient ineligible for IV tPA?

- No findings on head CT
- Stroke onset 4 hrs
- ASPECTS score of 2
- Positive basal ganglia sing on CT


34

# Acute Stroke – Imaging Evaluation




35

## CTA: Large Vessel Occlusion



36


## Collaterals



- **Good collaterals:** Smaller baseline infarct core, reduced final infarct volumes and larger mismatches at triage
- **Poor collaterals:** Larger core and quicker infarct growth
- Amount of **core** growth and **speed** of **growth** may depend on collaterals


37

## Acute Stroke – Imaging Evaluation



38

## CT Perfusion




- CT perfusion (CTP) has emerged as a valuable tool in the management of **acute stroke**, providing essential **quantitative** information on the extent of the irreversibly injured ischemic core and the potentially salvageable ischemic penumbra

39


## Salvageable Brain?

- Core:** Irreversible injured tissue
- Penumbra:** at risk, but salvageable, if reperfusion occurs




CTP help in the **selection of patients** for intravascular thrombolysis and endovascular revascularization

40



41



42

## Multiple Perfusion Parameters

### Tissue viability:


- CBF: Cerebral blood flow (nml 50-60ml/100gr/min)
- CBV: Cerebral blood volume (2-5mL/100g tissue)
- $CBF = CBV / MTT$

### Time-sensitive:


- MTT: Mean transit time (3-5 sec, needs deconvolution)
- TTP: Time to peak (maximum contrast variation)
- $TTP = T_{max}$ : Time to max (sec, needs deconvolution)
- TA: Time of arrival of the contrast agent

43


## CT Perfusion



44



45



46



## Examples CTP


47

## Case 1

88 yo F

Acute onset LUE weakness, L facial  
droop and dysarthria


48




49




50




51




52



53



54




55


## Case 2

63 yo F  
Code Stroke


56




57




58




59



60




61




62





63




64


## Ischemic Stroke

- Early hyperacute: 0 to 6 hours
- Late hyperacute: 6 to 24 hours
- Acute: 24 hours to 1 week
- Subacute: 1 to 3 weeks
- Chronic: more than 3 weeks


65



66




67



68

# Acute Stroke Imaging



## Jose Gavito-Higuera, MD

Assistant Professor, Neuroradiology Division  
Texas Tech University Health Sciences Center  
University Medical Center of El Paso  
El Paso, Texas  
[jose.gavito@ttuhsc.edu](mailto:jose.gavito@ttuhsc.edu)